



Dr. Ravish I.R.


MS, MCh. (Urology)
Fellowship in Paediatric Urology (KLEU)
Fellowship in Paediatric Urology (Minnesota, USA)
Consultant Urologist, Laparoscopic,
Endoscopic & Paediatric Urologist
email: irravish@yahoo.com

Dr. Ravish I. R. - He did his MBBS & MS (General Surgery) at JSS Medical College Mysore and MCh (Urology) from JNM Medical College Belgaum in the year 2007, followed by Fellowship in Paediatric Urology from KLES, Belgaum and later did his Fellow in Paediatric Urology from University of Minnesota (USA) in the year 2008. He presently has formed a formidable team to deliver the best in Paediatric Urology. He is the Sr. Urologist at Vikram Hospital, Bengaluru which consists a team of Urologist & Nephrologist.

Special Interest in:

- Paediatric Urology
- Laser Urology
- Laparoscopy

not for business solicitation for information only.



**Centre of excellence in
tertiary Uro Care.**



34, 31st 'A' Cross, Jayanagar 7th Block, Bengaluru - 560 011.

Ph: 080-2665 0004 / 82776 74213

Email: irravish@yahoo.com



Circumcision

is a surgery to remove the foreskin from the penis.

Any



pain after surgery can be controlled by giving pain control analgesics, antibiotics and applying antiseptic ointment over the operated area for faster recovery.

Doing

circumcision is sometimes the choice based on religious, social or belief, besides medical reasons.



The

medical reason of circumcision is to minimize the infection of bladder or foreskin.



Bring

loose underwear and pants for baby boy to use after surgery.

Most boys will be able to return to their usual activities (like sports, going to school) within few days after circumcision.



Circumcision

is most often done on a baby boy just before he goes home from the hospital. One can also choose to do it at the later date.



Call doctor if,

- ◆ Bleeding occurs
- ◆ Baby boy not urinating for at least every 8 hours
- ◆ Pain not relieving with the prescribed medicine
- ◆ There is high temperature
- ◆ Increase in swelling or redness around the area
- ◆ There is pus formation.



Getting

circumcised helps in washing penis easily. Washing reduces odor, swelling and infections.

Also, chances of getting sexually transmitted disease (std) or urinary infection reduces to a large extent.



Circumcision

surgery is performed under general anesthesia or regional block.

The penis and surrounding skin will be carefully washed with antiseptic solution and the foreskin is then removed.

After which the area is covered with antiseptic ointment.



One needs a circumcision, if there is:

- ◆ An obstruction to the flow of urine
 Yes No
- ◆ Ballooning of the foreskin
 Yes No
- ◆ Thin stream of urination
 Yes No
- ◆ Recurrent urinary tract infection
 Yes No

If 'Yes' to any of the above, meet our **"Consultant Urologists"** today.